



EVIDENCE-BASED
PRACTICES

KIT

Knowledge Informing Transformation

The Evidence

Integrated Treatment for Co-Occurring Disorders



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Substance Abuse and Mental Health Services Administration
Center for Mental Health Services
www.samhsa.gov

The Evidence



Integrated Treatment for Co-Occurring Disorders

Acknowledgments

This document was produced for the Substance Abuse and Mental Health Services Administration (SAMHSA) by the New Hampshire-Dartmouth Psychiatric Research Center under contract number 280-00-8049 and Westat under contract number 270-03-6005, with SAMHSA, U.S. Department of Health and Human Services (HHS). Neal Brown, M.P.A., and Crystal Blyler, Ph.D., served as the Government Project Officers.

Disclaimer

The views, opinions, and content of this publication are those of the authors and contributors and do not necessarily reflect the views, opinions, or policies of the Center for Mental Health Services (CMHS), SAMHSA, or HHS.

Public Domain Notice

All material appearing in this document is in the public domain and may be reproduced or copied without permission from SAMHSA. Citation of the source is appreciated. However, this publication may not be reproduced or distributed for a fee without the specific, written authorization from the Office of Communications, SAMHSA, HHS.

Electronic Access and Copies of Publication

This publication may be downloaded or ordered at www.samhsa.gov/shin. Or, please call SAMHSA's Health Information Network at **1-877-SAMHSA-7** (1-877-726-4727) (English and Español).

Recommended Citation

Substance Abuse and Mental Health Services Administration. *Integrated Treatment for Co-Occurring Disorders: The Evidence*. DHHS Pub. No. SMA-08-4366, Rockville, MD: Center for Mental Health Services, Substance Abuse and Mental Health Services Administration, U.S. Department of Health and Human Services, 2009.

Originating Office

Center for Mental Health Services
Substance Abuse and Mental Health Services Administration
1 Choke Cherry Road
Rockville, MD 20857

DHHS Publication No. SMA-08-4366
Printed 2009



The Evidence

The Evidence introduces all stakeholders to the research literature and other resources on Integrated Treatment for Co-Occurring Disorders. This booklet includes the following:

- Two key resources included in the KIT;
- Additional resources for further reading; and
- References for the citations presented throughout the KIT.

Integrated Treatment for Co-Occurring Disorders

This KIT is part of a series of Evidence-Based Practices KITs created by the Center for Mental Health Services, Substance Abuse and Mental Health Services Administration, U.S. Department of Health and Human Services.

This booklet is part of the Integrated Treatment for Co-Occurring Disorders KIT that includes a DVD, CD-ROM, and seven booklets:

How to Use the Evidence-Based Practices KITs

Getting Started with Evidence-Based Practices

Building Your Program

Training Frontline Staff

Evaluating Your Program

The Evidence

Using Multimedia to Introduce Your EBP



EVIDENCE-BASED
PRACTICES

KIT

Knowledge Informing Transformation

What's in *The Evidence*

Review of Research Literature	1
Selected Bibliography	11
References	23
Acknowledgments	27

Integrated Treatment for Co-Occurring Disorders

The Evidence

Review of Research Literature

A number of research articles summarize the effectiveness of Integrated Treatment for Co-Occurring Disorders. This KIT includes a full text copy of one of them:

Drake, R. E., Essock, S. M., Shaner, A., Carey, K. B., Minkoff, K., Kola, L., et al. (2001). Implementing dual diagnosis services for clients with severe mental illness. *Psychiatric Services*, 52(4), 469-476.

This article describes the critical components of the evidence-based model and its effectiveness. Barriers to implementation and strategies for overcoming them are also discussed, based on experiences in several states.

This article may be viewed or printed from the CD-ROM in your KIT. For a printed copy, see page 3.



Center for Substance Abuse Treatment, Substance Abuse Treatment for Persons with Co-Occurring Disorders, Treatment Improvement Protocol (TIP) Series 42, DHHS Publication No. (SMA) 05-3992, Rockville, MD: Substance Abuse and Mental Health Services Administration, 2005.

Similar to this KIT, TIP 42 is a guide that provides resources for treating co-occurring mental illnesses and substance use disorders. It is an excellent complement to the Integrated Treatment KIT.

The primary audiences for TIP 42 are substance abuse treatment practitioners with varying degrees of education and experience. Secondary audiences are policymakers and other professionals who work with consumers who have co-occurring disorders.

TIP 42 summarizes state-of-the-art treatment of co-occurring disorders. It has chapters on terminology, assessment, and treatment strategies and gives suggestions for policy planning.

Concepts, models, and strategies outlined in TIP 42 are based on definitive research, empirical support, and agreements of a consensus panel. Successful models of treatment are portrayed and specific consensus panel recommendations are cited throughout the TIP.

TIP 42 is accompanied by four supplements:

- *Quick Guide for Administrators Based on TIP 42: Substance Abuse Treatment for Persons with Co-Occurring Disorders;*
- *Quick Guide for Clinicians Based on TIP 42: Substance Abuse Treatment for Persons with Co-Occurring Disorders;*
- *Substance Abuse Treatment for Persons with Co-Occurring Disorders: In-Service Training Based on a Treatment Improvement Protocol Tip 42; and*
- *Knowledge Application Program (KAP) Keys for Clinicians Based on TIP 42: Substance Abuse Treatment for Persons with Co-Occurring Disorders.*

For a copy of TIP 42 and the supplemental guides for the TIP, see the CD-ROM in this KIT or go to www.ncadi.samhsa.gov.

Implementing Dual Diagnosis Services for Clients With Severe Mental Illness

2001
Dedicated to
Evidence-
Based
Psychiatry

Robert E. Drake, M.D., Ph.D.
Susan M. Essock, Ph.D.
Andrew Shaner, M.D.
Kate B. Carey, Ph.D.
Kenneth Minkoff, M.D.
Lenore Kola, Ph.D.
David Lynde, M.S.W.
Fred C. Osher, M.D.
Robin E. Clark, Ph.D.
Lawrence Rickards, Ph.D.

After 20 years of development and research, dual diagnosis services for clients with severe mental illness are emerging as an evidence-based practice. Effective dual diagnosis programs combine mental health and substance abuse interventions that are tailored for the complex needs of clients with comorbid disorders. The authors describe the critical components of effective programs, which include a comprehensive, long-term, staged approach to recovery; assertive outreach; motivational interventions; provision of help to clients in acquiring skills and supports to manage both illnesses and to pursue functional goals; and cultural sensitivity and competence. Many state mental health systems are implementing dual diagnosis services, but high-quality services are rare. The authors provide an overview of the numerous barriers to implementation and describe implementation strategies to overcome the barriers. Current approaches to implementing dual diagnosis programs involve organizational and financing changes at the policy level, clarity of program mission with structural changes to support dual diagnosis services, training and supervision for clinicians, and dissemination of accurate information to consumers and families to support understanding, demand, and advocacy. (*Psychiatric Services* 52:469–476, 2001)

Substance abuse is the most common and clinically significant comorbid disorder among adults with severe mental illness. In this paper the term “substance abuse” refers to substance use disorders, which include abuse and dependence. “Severe mental illness” refers to long-term psychiatric disorders, such as schizophrenia, that are associated with disability and that fall within the traditional purview of public mental health systems. Finally, the term “dual diagnosis” denotes the co-occurrence of substance abuse and severe mental illness.

There are many populations with dual diagnoses, and there are other common terms for this particular group. Furthermore, dual diagnosis is a misleading term because the individuals in this group are heterogeneous and tend to have multiple impairments rather than just two illnesses. Nevertheless, the term appears consistently in the literature and has acquired some coherence as a referent to particular clients, treatments, programs, and service system issues.

Since the problem of dual diagnosis became clinically apparent in the early 1980s (1,2), researchers have established three basic and consistent findings. First, co-occurrence is common;

Dr. Drake and Dr. Clark are affiliated with the New Hampshire–Dartmouth Psychiatric Research Center, 2 Whipple Place, Lebanon, New Hampshire 03766 (e-mail, robert.e.drake@dartmouth.edu). Dr. Essock is with Mt. Sinai Medical School in New York City. Dr. Shaner is affiliated with the School of Medicine at the University of California, Los Angeles. Dr. Carey is with Syracuse University in New York. Dr. Minkoff is in private practice in Boston. Dr. Kola is with Case Western Reserve University in Cleveland, Ohio. Mr. Lynde is with West Institute in Concord, New Hampshire. Dr. Osher is with the University of Maryland School of Medicine in Baltimore. Dr. Rickards is with the Center for Mental Health Services in Rockville, Maryland.

PSYCHIATRIC SERVICES April 2001 Vol. 52 No. 4

469

about 50 percent of individuals with severe mental disorders are affected by substance abuse (3). Second, dual diagnosis is associated with a variety of negative outcomes, including higher rates of relapse (4), hospitalization (5), violence (6), incarceration (7), homelessness (8), and serious infections such as HIV and hepatitis (9). Third, the parallel but separate mental health and substance abuse treatment systems so common in the United States deliver fragmented and ineffective care (10). Most clients are unable to navigate the separate systems or make sense of disparate messages about treatment and recovery. Often they are excluded or extruded from services in one system because of the comorbid disorder and told to return when the other problem is under control. For those reasons, clinicians, administrators, researchers, family organizations, and clients themselves have been calling for the integration of mental health and substance abuse services for at least 15 years (10,11).

Over that time, integrated dual diagnosis services—that is, treatments and programs—have been steadily developed, refined, and evaluated (11). This paper, part of a series on specific evidence-based practices for persons with severe mental illness, provides an overview of the evolution of dual diagnosis services, the evidence on outcomes and critical components, and the limitations of current research. We also address barriers to the implementation of dual diagnosis services and current strategies for implementation in routine mental health settings.

Dual diagnosis services

Treatments, or interventions, are offered within programs that are part of service systems. Dual diagnosis treatments combine or integrate mental health and substance abuse interventions at the level of the clinical interaction. Hence integrated treatment means that the same clinicians or teams of clinicians, working in one setting, provide appropriate mental health and substance abuse interventions in a coordinated fashion. In other words, the caregivers take responsibility for combining the interventions

Editor's note: This article is part of a series of papers on evidence-based practices being published in *Psychiatric Services* this year. The papers focus on mental health practices for which there is substantial evidence of effectiveness and that should therefore be routinely offered in clinical settings. Articles in previous issues have addressed implementing evidence-based practices for persons with severe mental illness and in routine mental health settings and implementing supported employment as an evidence-based practice. Among other topics, future articles will examine evidence-based practices for case management and assertive community treatment, illness self-management, children's services, and services for the elderly population. Robert E. Drake, M.D., Ph.D., and Howard H. Goldman, M.D., Ph.D., are the series editors.

into one coherent package. For the individual with a dual diagnosis, the services appear seamless, with a consistent approach, philosophy, and set of recommendations. The need to negotiate with separate clinical teams, programs, or systems disappears.

Integration involves not only combining appropriate treatments for both disorders but also modifying traditional interventions (12–15). For example, social skills training emphasizes the importance of developing relationships but also the need to avoid social situations that could lead to substance use. Substance abuse counseling goes slowly, in accordance with the cognitive deficits, negative symptoms, vulnerability to confrontation, and greater need for support that are characteristic of many individuals with severe mental illness. Family interventions address understanding and learning to cope with two interacting illnesses.

The goal of dual diagnosis interventions is recovery from two serious ill-

nesses (16). In this context, “recovery” means that the individual with a dual diagnosis learns to manage both illnesses so that he or she can pursue meaningful life goals (17,18).

Research on dual diagnosis practices

In most states, the publicly financed mental health system bears responsibility for providing treatments and support services for clients with severe mental illness. Dual diagnosis treatments for these clients have therefore generally been added to community support programs within the mental health system.

Early studies of dual diagnosis interventions during the 1980s examined the application of traditional substance abuse treatments, such as 12-step groups, to clients with mental disorders within mental health programs. These studies had disappointing results for at least two reasons (19). The clinical programs did not take into account the complex needs of the population, and researchers had not yet solved basic methodologic problems. For example, early programs often failed to incorporate outreach and motivational interventions, and evaluations were limited by lack of reliable and valid assessment of substance abuse. Reviews based on these early studies were understandably pessimistic (20).

At the same time, however, a series of demonstration projects using more comprehensive programs that incorporated assertive outreach and long-term rehabilitation began to show better outcomes. Moreover, the projects developed motivational interventions to help clients who did not perceive or acknowledge their substance abuse or mental illness problems (21).

Building on these insights, projects in the early 1990s incorporated motivational approaches as well as outreach, comprehensiveness, and a long-term perspective, often within the structure of multidisciplinary treatment teams. These later studies, which were uncontrolled but incorporated more valid measures of substance abuse, generally showed positive outcomes, including substantial rates of stable remission of substance abuse (22–25). Of course, uncon-

trolled studies of this type often produce findings that are not replicated in controlled studies; they should be considered pilot studies, which are often needed to refine the intervention and the methodologies of evaluation and which should be followed by controlled investigation to determine evidence-based practice (26).

Controlled research studies of comprehensive dual diagnosis programs began to appear in the mid-1990s. Eight recent studies with experimental or quasi-experimental designs support the effectiveness of integrated dual diagnosis treatments for clients with severe mental illness and substance use disorders (27–34). The type and array of dual diagnosis interventions in these programs vary, but they include several common components, which are reviewed below. The eight studies demonstrated a variety of positive outcomes in domains such as substance abuse, psychiatric symptoms, housing, hospitalization, arrests, functional status, and quality of life (19). Although each had methodological limitations, together they indicate that current integrated treatment programs are more effective than nonintegrated programs. By contrast, the evidence continues to show that dual diagnosis clients in mental health programs that fail to integrate substance abuse interventions have poor outcomes (35).

Critical components

Several components of integrated programs can be considered evidence-based practices because they are almost always present in programs that have demonstrated good outcomes in controlled studies and because their absence is associated with predictable failures (21). For example, dual diagnosis programs that include assertive outreach are able to engage and retain clients at a high rate, while those that fail to include outreach lose many clients.

Staged interventions

Effective programs incorporate, implicitly or explicitly, the concept of stages of treatment (14,36,37). In the simplest conceptualization, stages of treatment include forming a trusting

relationship (engagement), helping the engaged client develop the motivation to become involved in recovery-oriented interventions (persuasion), helping the motivated client acquire skills and supports for controlling illnesses and pursuing goals (active treatment), and helping the client in stable remission develop and use strategies for maintaining recovery (relapse prevention).

Clients do not move linearly through stages. They sometimes enter services at advanced levels, skip over or pass rapidly through stages, or relapse to earlier stages. They may be in different stages with respect to mental illness and substance abuse. Nevertheless, the concept of stages has proved useful to program planners and clinicians because clients at different stages respond to stage-specific interventions.

Assertive outreach

Many clients with a dual diagnosis have difficulty linking with services and participating in treatment (38). Effective programs engage clients and members of their support systems by providing assertive outreach, usually through some combination of intensive case management and meetings in the client's residence (21,32). For example, homeless persons with dual diagnoses often benefit from outreach, help with housing, and time to develop a trusting relationship before participating in any formal treatment. These approaches enable clients to gain access to services and maintain needed relationships with a consistent program over months and years. Without such efforts, noncompliance and dropout rates are high (39).

Motivational interventions

Most dual diagnosis clients have little readiness for abstinence-oriented treatment (40,41). Many also lack motivation to manage psychiatric illness and to pursue employment or other functional goals. Effective programs therefore incorporate motivational interventions that are designed to help clients become ready for more definitive interventions aimed at illness self-management (12,14,21). For example, clients who

are so demoralized, symptomatic, or confused that they mistakenly believe that alcohol and cocaine are helping them to cope better than medications require education, support, and counseling to develop hope and a realistic understanding of illnesses, drugs, treatments, and goals.

Motivational interventions involve helping the individual identify his or her own goals and to recognize, through a systematic examination of the individual's ambivalence, that not managing one's illnesses interferes with attaining those goals (42). Recent research has demonstrated that clients who are not motivated can be reliably identified (43) and effectively helped with motivational interventions (Carey KB, Carey MP, Maisto SA, et al, unpublished data, 2000).

Counseling

Once clients are motivated to manage their own illnesses, they need to develop skills and supports to control symptoms and to pursue an abstinent lifestyle. Effective programs provide some form of counseling that promotes cognitive and behavioral skills at this stage. The counseling takes different forms and formats, such as group, individual, or family therapy or a combination (15). Few studies have compared specific approaches to counseling, although one study did find preliminary evidence that a cognitive-behavioral approach was superior to a 12-step approach (28). At least three research groups are actively working to refine cognitive-behavioral approaches to substance abuse counseling for dual diagnosis clients (12,13,44). These approaches often incorporate motivational sessions at the beginning of counseling and as needed in subsequent sessions rather than as separate interventions.

Social support interventions

In addition to helping clients build skills for managing their illness and pursuing goals, effective programs focus on strengthening the immediate social environment to help them modify their behavior. These activities, which recognize the role of social networks in recovery from dual disorders (45), include social network or family interventions.

Long-term perspective

Effective programs recognize that recovery tends to occur over months or years in the community. People with severe mental illness and substance abuse do not usually develop stability and functional improvements quickly, even in intensive treatment programs, unless they enter treatment at an advanced stage (19). Instead, they tend to improve over months and years in conjunction with a consistent dual diagnosis program. Effective programs therefore take a long-term, community-based perspective that includes rehabilitation activities to prevent relapses and to enhance gains.

Comprehensiveness

Learning to lead a symptom-free, abstinent lifestyle that is satisfying and sustainable often requires transforming many aspects of one's life—for example, habits, stress management, friends, activities, and housing. Therefore, in effective programs attention to substance abuse as well as mental illness is integrated into all aspects of the existing mental health program and service system rather than isolated as a discrete substance abuse treatment intervention. Inpatient hospitalization, assessment, crisis intervention, medication management, money management, laboratory screening, housing, and vocational rehabilitation incorporate special features that are tailored specifically for dual diagnosis patients. For example, hospitalization is considered a component of the system that supports movement toward recovery by providing diagnosis, stabilization, and linkage with outpatient dual diagnosis interventions during acute episodes (46). Similarly, housing and vocational programs can be used to support the individual with a dual diagnosis in acquiring skills and supports needed for recovery (47).

Cultural sensitivity and competence

A fundamental finding of the demonstration programs of the late 1980s was that cultural sensitivity and competence were critical to engaging clients in dual diagnosis services (21). These demonstrations showed that

African Americans, Hispanics, and other underserved groups, such as farm workers, homeless persons, women with children, inner-city residents, and persons in rural areas, could be engaged in dual diagnosis services if the services were tailored to their particular racial, cultural, and other group characteristics.

Many dual diagnosis programs omit some of these critical components as evidence-based practices. However, one consistent finding in the research is that programs that show high fidelity to the model described here—those that incorporate more of the core elements—produce better outcomes than low-fidelity programs (32,48,49). A common misconception about technology transfer is that model programs are not generalizable and that local solutions are superior. A more accurate reading of the research is that modifications for cultural and other local circumstances are important, but critical program components must be replicated to achieve good outcomes.

Limitations of the research

The design and quality of research procedures and data across dual diagnosis studies are inconsistent. In addition, researchers have thus far failed to address a number of issues.

Dual diagnosis research has studied the clinical enterprise, that is, treatments and programs, with little attention to the policy or system perspective. Despite widespread endorsement of integrated dual diagnosis services (13,50–53), there continues to be a general failure at the federal and state levels to resolve problems related to organization and financing (see below). Thus, despite the emergence of many excellent programs around the country, few if any large mental health systems have been able to accomplish widespread implementation of dual diagnosis services for persons with severe mental illness. We are aware of no specific studies of strategies to finance, contract for, reorganize, or train in relation to dual diagnosis services.

Lack of data on the cost of integrated dual diagnosis services and the cost savings of providing good care impedes policy development. Dual

diagnosis clients incur high treatment costs in usual services (54,55), and care is costly to their families (56), but effective treatment may be even more costly. Some studies suggest cost savings related to providing good services (57,58), but these are not definitive.

Another limitation of the research is the lack of specificity of dual diagnosis treatments. Interventions differ across studies, manuals and fidelity measures are rare, and no consensus exists on specific approaches to individual counseling, group treatment, family intervention, housing, medications, and other components. Current research will address some of these issues by refining specific components, although efficacy studies may identify complex and expensive interventions that will be impractical in routine mental health settings.

A majority of dual diagnosis clients respond well to integrated outpatient services, but clients who do not respond continue to be at high risk of hospitalization, incarceration, homelessness, HIV infection, and other serious adverse outcomes. Other than one study of long-term residential treatment (33), controlled research has not addressed clients who do not respond to outpatient services. Other potential interventions include outpatient commitment (59), treatments aimed at trauma sequelae (60), money management (61), contingency management (62), and pharmacological approaches using medications such as clozapine (63), disulfiram (64), or naltrexone.

Although a few studies have explored the specific treatment needs of dual diagnosis clients who are women (65,66) or minorities (21,67), particular program modifications for these groups need further validation. For example, many dual diagnosis programs have identified high rates of trauma histories and sequelae among women (46,68,69), and studies have suggested interventions to address trauma; however, no data on outcomes are yet available.

Implementation barriers

Although integrated dual diagnosis services and other evidence-based practices are widely advocated, they

are rarely offered in routine mental health treatment settings (70). The barriers are legion.

Policy barriers

State, county, and city mental health authorities often encounter policies related to organizational structure, financing, regulations, and licensing that militate against the functional integration of mental health and substance abuse services (71). The U.S. public mental health and substance abuse treatment systems grew independently. In most states these services are provided under the auspices of separate cabinet-level departments with separate funding streams, advocacy groups, lobbyists, enabling legislation, information systems, job classifications, and criteria for credentials. Huge fiscal incentives and strong political allies act to maintain the status quo.

Medicaid programs, which fund a significant and growing proportion of treatment for persons with severe mental illness, vary substantially from state to state in the types of mental health and substance abuse services they fund. In most states, mental health and substance abuse agencies have little control over how Medicaid services are reimbursed or administered, which makes it difficult for public systems to ensure that appropriate services are accessible. Medicare, the federal insurance program for elderly and disabled persons, generally pays for a more limited scope of mental health and substance abuse services. Together Medicaid and Medicare pay for more than 30 percent of all behavioral health services, but their impact on dual diagnosis services has not been studied (72).

Program barriers

At the local level, administrators of clinics, centers, and programs have often lacked the clear service models, administrative guidelines, contractual incentives, quality assurance procedures, and outcome measures needed to implement dual diagnosis services. When clinical needs compel them to move ahead anyway, they have difficulty hiring a skilled workforce with experience in provid-

ing dual diagnosis interventions and lack the resources to train current supervisors and clinicians.

Clinical barriers

The beliefs of the mental health and substance abuse treatment traditions are inculcated in clinicians, which diminishes the opportunities for cross-fertilization (73). Although an integrated clinical philosophy and a practical approach to dual diagnosis treatment have been clearly delineated for more than a decade (16), educational institutions rarely teach this approach. Consequently, mental health clinicians typically lack training in dual diagnosis treatment and have to rely on informal, self-initiated opportunities for learning current interventions (74). They often avoid diagnosing substance abuse when they believe that it is irrelevant, that it will interfere with funding, or that they cannot treat it. Clinicians trained in substance abuse treatment, as well as recovering dual diagnosis clients, could add expertise and training, but they are often excluded from jobs in the mental health system.

Consumer and family barriers

Clients and their families rarely have good information about dual diagnosis and appropriate services. Few programs offer psychoeducational services related to dual diagnosis, although practical help from families plays a critical role in recovery (75). Family members are often unaware of substance abuse, blame all symptoms on drug abuse, or attribute symptoms and substance use to willful misbehavior. Supporting family involvement is an important but neglected role for clinicians.

Consumers often deny or minimize problems related to substance abuse (40) and, like other substance abusers, believe that alcohol or other drugs are helpful in alleviating distress. They may be legitimately confused about causality because they perceive the immediate effects of drugs rather than the intermediate or long-term consequences (76). The net result is that the individual lacks motivation to pursue active substance abuse treatment, which can reinforce clinical inattention.

Implementation strategies

There are no proven strategies for overcoming the aforementioned barriers to implementing dual diagnosis services, but some suggestions have come from systems and programs that have had moderate success.

Policy strategies

Health care authorities in a majority of, and possibly all, states have current initiatives for creating dual diagnosis services. Because health care policy is often administered at the county or city level, hundreds of individual experiments are occurring. One initial branch point involves the decision to focus broadly on the entire behavioral health system—that is, on all clients with mental health and substance abuse problems—or more narrowly on services for those with severe mental illness and co-occurring substance abuse. We examine here only strategies for dual diagnosis clients with severe mental illness, for whom the implementation issues are relatively distinct.

Commonly used system-level strategies include building a consensus around the vision for integrated services and then conjointly planning; specifying a model; implementing structural, regulatory, and reimbursement changes; establishing contracting mechanisms; defining standards; and funding demonstration programs and training initiatives (77). To our knowledge, few efforts have been made to study these efforts at the system level.

Anecdotal evidence indicates that blending mental health and substance abuse funds appears to have been a relatively unsuccessful strategy, especially early in the course of system change. Fear of losing money to cover nontraditional populations often leads to prolonged disagreements, inability to develop consensus, and abandonment of other plans. As a less controversial, preliminary step, the mental health authority often assumes responsibility for comprehensive care, including substance abuse treatment, for persons with severe mental illness, while the substance abuse authority assists by pledging to help with training and planning.

This limited approach enables the

mental health system to attract and train dual diagnosis specialists who can subsequently train other clinicians and programs. Without structural, regulatory, and funding changes to reinforce the training, however, the expertise may soon disappear—a common experience after demonstration projects. Thus many experts advise that policy issues should be addressed early in the process of implementation to avoid wasting efforts on training (78–80).

New costs to the mental health system for dual diagnosis training could be offset by greater effectiveness in ameliorating substance-abusing behaviors that are associated with hospitalizations. However, saving costs over time assumes that providers are at risk for all treatment costs, that is, that providers have incentives to invest more in outpatient services in order to spend less on inpatient services. Despite the growth of managed care, providers rarely bear complete financial responsibility for the treatment of clients with severe mental illness.

Program strategies

At the level of the mental health clinic or program leadership, the fundamental task is to begin recognizing and treating substance abuse rather than ignoring it or using it as a criterion for exclusion (81). After consensus-building activities to prepare for change, staff need training and supervision to learn new skills, and they must receive reinforcement for acquiring and using these skills effectively. One common strategy is to appoint a director of dual diagnosis services whose job is to plan and oversee the training of staff, the integration of substance abuse awareness and treatment into all aspects of the mental health program, and the monitoring and reinforcement of these activities through medical records, quality assurance activities, and outcome data.

Experts identify the importance of having a single leader for program change (82). Fidelity measures for integrated dual diagnosis services can facilitate successful implementation at the program level (50,83). Monitoring and reinforcing mechanisms

also emphasize client-centered outcomes, such as abstinence and employment.

Clinical strategies

Mental health clinicians need to acquire knowledge and a core set of skills related to substance abuse that includes assessing substance abuse, providing motivational interventions for clients who are not ready to participate in abstinence-oriented treatment, and providing counseling for those who are motivated try to maintain abstinence. Clinicians adopt new skills as a result of motivation, instruc-

tion, practice, and reinforcement (84). Because substance abuse affects the lives of the great majority of clients with severe mental illness—as a co-occurring disorder, family stressor, or environmental hazard—all clinicians should learn these basic skills. Otherwise substance abuse problems will continue to be missed and untreated in this population (85,86).

tion, practice, and reinforcement (84). Because substance abuse affects the lives of the great majority of clients with severe mental illness—as a co-occurring disorder, family stressor, or environmental hazard—all clinicians should learn these basic skills. Otherwise substance abuse problems will continue to be missed and untreated in this population (85,86). For example, all case managers should recognize and address substance abuse in their daily interactions, as should housing staff, employ-

Consumer- and family-level strategies

Clients and family members need access to accurate information. Otherwise their opportunities to make informed choices, to request effective services, and to advocate for system changes are severely compromised. Consumer demand and family advocacy can move the health care system toward evidence-based practices, but concerted efforts at the national, state, and local levels are required. Researchers can facilitate their efforts by offering clear messages about the forms, processes, and expected outcomes of evidence-based practices. Similarly, local programs should provide information on available dual diagnosis services to clients and their families.

As consumers move into roles as providers within the mental health system and in consumer-run services, they also need training in dual diagnosis treatments. Local educational programs, such as community colleges, as well as staff training programs should address these needs.

Conclusions

Substance abuse is a common and devastating comorbid disorder among persons with severe mental illness. Recent research offers evidence that integrated dual diagnosis treatments are effective, but basic interventions are rarely incorporated into the mental health programs in which these clients receive care. Successful implementation of dual diagnosis services within mental health systems will depend on changes at several levels: clear policy directives with consistent organizational and financing supports, program changes to incorpo-

■

*Recent
research
offers evidence
that integrated dual
diagnosis treatments
are effective, but basic
interventions are rarely
incorporated into the
mental health programs
in which these
clients receive
care.*

■

rate the mission of addressing co-occurring substance abuse, supports for the acquisition of expertise at the clinical level, and availability of accurate information to consumers and family members.

Acknowledgments

This review by a national panel was supported by grant 036805 from the Robert Wood Johnson Foundation and contract 280-00-8049 from the Center for Mental Health Services.

References

1. Pepper B, Krishner MC, Ryglewicz H: The young adult chronic patient: overview of a population. *Hospital and Community Psychiatry* 32:463–469, 1981
2. Caton CLM: The new chronic patient and the system of community care. *Hospital and Community Psychiatry* 32:475–478, 1981
3. Regier DA, Farmer ME, Rae DS, et al: Comorbidity of mental disorders with alcohol and other drug abuse. *JAMA* 264:2511–2518, 1990
4. Swofford C, Kasckow J, Scheller-Gilkey G, et al: Substance use: a powerful predictor of relapse in schizophrenia. *Schizophrenia Research* 20:145–151, 1996
5. Haywood TW, Kravitz HM, Grossman LS, et al: Predicting the “revolving door” phenomenon among patients with schizophrenic, schizoaffective, and affective disorders. *American Journal of Psychiatry* 152:856–861, 1995
6. Cuffel B, Shumway M, Chouljian T: A longitudinal study of substance use and community violence in schizophrenia. *Journal of Nervous and Mental Disease* 182:704–708, 1994
7. Abram KM, Teplin LA: Co-occurring disorders among mentally ill jail detainees: implications for public policy. *American Psychologist* 46:1036–1045, 1991
8. Caton CLM, Shrout PE, Eagle PF, et al: Risk factors for homelessness among schizophrenic men: a case-control study. *American Journal of Public Health* 84:265–270, 1994
9. Rosenberg SD, Goodman LA, Osher FC, et al: Prevalence of HIV, hepatitis B, and hepatitis C in people with severe mental illness. *American Journal of Public Health* 91:31–37, 2001
10. Ridgely MS, Osher FC, Goldman HH, et al: Executive Summary: Chronic Mentally Ill Young Adults With Substance Abuse Problems: A Review of Research, Treatment, and Training Issues. Baltimore, University of Maryland School of Medicine, Mental Health Services Research Center, 1987
11. Drake RE, Wallach MA: Dual diagnosis: 15 years of progress. *Psychiatric Services* 51:1126–1129, 2000
12. Barrowclough C, Haddock G, Tarrier N, et al: Cognitive-behavioral intervention for clients with severe mental illness who have a substance misuse problem. *Psychiatric Rehabilitation Skills* 4:216–233, 2000
13. Bellack AS, DiClemente CC: Treating substance abuse among patients with schizophrenia. *Psychiatric Services* 50:75–80, 1999
14. Carey KB: Substance use reduction in the context of outpatient psychiatric treatment: a collaborative, motivational, harm reduction approach. *Community Mental Health Journal* 32:291–306, 1996
15. Mueser KT, Drake RE, Noordsy DL: Integrated mental health and substance abuse treatment for severe psychiatric disorders. *Journal of Practical Psychiatry and Behavioral Health* 4:129–139, 1998
16. Minkoff K: An integrated treatment model for dual diagnosis of psychosis and addiction. *Hospital and Community Psychiatry* 40:1031–1036, 1989
17. Mead S, Copeland ME: What recovery means to us: consumers’ perspectives. *Community Mental Health Journal* 36:315–328, 2000
18. Torrey WC, Wyzik P: The recovery vision as a service improvement guide for community mental health center providers. *Community Mental Health Journal* 36:209–216, 2000
19. Drake RE, Mercer-McFadden C, Mueser KT, et al: Review of integrated mental health and substance abuse treatment for patients with dual disorders. *Schizophrenia Bulletin* 24:589–608, 1998
20. Ley A, Jeffery DP, McLaren S, et al: Treatment programmes for people with both severe mental illness and substance misuse. *Cochrane Library*, Feb 1999. Available at <http://www.update-software.com/cochrane/cochrane-frame.html>
21. Mercer-McFadden C, Drake RE, Brown NB, et al: The community support program demonstrations of services for young adults with severe mental illness and substance use disorders, 1987–1991. *Psychiatric Rehabilitation Journal* 20(3):13–24, 1997
22. Detrick A, Stieppock V: Treating persons with mental illness, substance abuse, and legal problems: the Rhode Island experience. *New Directions for Mental Health Services*, no 56:65–77, 1992
23. Drake RE, McHugo G, Noordsy DL: Treatment of alcoholism among schizophrenic outpatients: four-year outcomes. *American Journal of Psychiatry* 150:328–329, 1993
24. Durell J, Lechtenberg B, Corse S, et al: Intensive case management of persons with chronic mental illness who abuse substances. *Hospital and Community Psychiatry* 44:415–416, 1993
25. Meisler N, Blankertz L, Santos AB, et al: Impact of assertive community treatment on homeless persons with co-occurring severe psychiatric and substance use disorders. *Community Mental Health Journal* 33:113–122, 1997
26. Drake RE, Goldman HH, Leff HS, et al: Implementing evidence-based practices in routine mental health service settings. *Psychiatric Services* 52:179–182, 2001
27. Godley SH, Hoewing-Roberson R, Godley MD: Final MISA Report. Bloomington, Ill, Lighthouse Institute, 1994
28. Jerrell JM, Ridgely MS: Comparative effectiveness of three approaches to serving people with severe mental illness and substance abuse disorders. *Journal of Nervous and Mental Disease* 183:566–576, 1995
29. Drake RE, Yovetich NA, Bebout RR, et al: Integrated treatment for dually diagnosed homeless adults. *Journal of Nervous and Mental Disease* 185:298–305, 1997
30. Carnichael D, Tackett-Gibson M, Dell O, et al: Texas Dual Diagnosis Project Evaluation Report, 1997–1998. College Station, Tex, Texas A&M University, Public Policy Research Institute, 1998
31. Drake RE, McHugo GJ, Clark RE, et al: Assertive community treatment for patients with co-occurring severe mental illness and substance use disorder: a clinical trial. *American Journal of Orthopsychiatry* 68:201–215, 1998
32. Ho AP, Tsuang JW, Liberman RP, et al: Achieving effective treatment of patients with chronic psychotic illness and comorbid substance dependence. *American Journal of Psychiatry* 156:1765–1770, 1999
33. Brunette MF, Drake RE, Woods M, et al: A comparison of long-term and short-term residential treatment programs for dual diagnosis patients. *Psychiatric Services* 52:526–528, 2001
34. Barrowclough C, Haddock G, Tarrier N, et al: Randomised controlled trial of motivational interviewing and cognitive behavioural intervention for schizophrenia patients with associated drug or alcohol misuse. *American Journal of Psychiatry*, in press
35. Havassy BE, Shopshire MS, Quigley LA: Effects of substance dependence on outcomes of patients in a randomized trial of two case management models. *Psychiatric Services* 51:639–644, 2000
36. Osher FC, Kofoed LL: Treatment of patients with psychiatric and psychoactive substance use disorders. *Hospital and Community Psychiatry* 40:1025–1030, 1989
37. McHugo GJ, Drake RE, Burton HL, et al: A scale for assessing the stage of substance abuse treatment in persons with severe mental illness. *Journal of Nervous and Mental Disease* 183:762–767, 1995
38. Owen C, Rutherford V, Jones M, et al: Non-compliance in psychiatric aftercare. *Community Mental Health Journal* 33:25–34, 1997
39. Hellerstein DJ, Rosenthal RN, Miner CR: A prospective study of integrated outpatient treatment for substance-abusing schizophrenic patients. *American Journal on Addictions* 42:33–42, 1995
40. Test MA, Wallish LS, Allness DG, et al: Substance use in young adults with schizophrenic disorders. *Schizophrenia Bulletin* 15:465–476, 1989
41. Ziedonis D, Trudeau K: Motivation to quit

- using substances among individuals with schizophrenia: implications for a motivation-based treatment model. *Schizophrenia Bulletin* 23:229–238, 1997
42. Miller W, Rollnick S: *Motivational Interviewing: Preparing People to Change Addictive Behavior*. New York, Guilford, 1991
 43. Carey KB, Maisto SA, Carey MP, et al: Measuring readiness-to-change substance misuse among psychiatric outpatients: I. reliability and validity of self-report measures. *Journal of Studies on Alcohol*, in press
 44. Roberts LJ, Shaner A, Eckman T: *Overcoming Addictions: Skills Training for People With Schizophrenia*. New York, Norton, 1999
 45. Alverson H, Alverson M, Drake RE: An ethnographic study of the longitudinal course of substance abuse among people with severe mental illness. *Community Mental Health Journal* 36:557–569, 2000
 46. Greenfield SF, Weiss RD, Tohen M: Substance abuse and the chronically mentally ill: a description of dual diagnosis treatment services in a psychiatric hospital. *Community Mental Health Journal* 31:265–278, 1995
 47. Drake RE, Mueser KT: Psychosocial approaches to dual diagnosis. *Schizophrenia Bulletin* 26:105–118, 2000
 48. Jerrell JM, Ridgely MS: Impact of robustness of program implementation on outcomes of clients in dual diagnosis programs. *Psychiatric Services* 50:109–112, 1999
 49. McHugo GJ, Drake RE, Teague GB, et al: Fidelity to assertive community treatment and client outcomes in the New Hampshire dual disorders study. *Psychiatric Services* 50:818–824, 1999
 50. Rach-Beisel J, Scott J, Dixon L: Co-occurring severe mental illness and substance use disorders: a review of recent research. *Psychiatric Services* 50:1427–1434, 1999
 51. Mueser KT, Bellack A, Blanchard J: Comorbidity of schizophrenia and substance abuse. *Journal of Consulting and Clinical Psychology* 60:845–856, 1992
 52. Osher FC, Drake RE: Reversing a history of unmet needs: approaches to care for persons with co-occurring addictive and mental disorders. *American Journal of Orthopsychiatry* 66:4–11, 1996
 53. Woody G: The challenge of dual diagnosis. *Alcohol Health and Research World* 20:76–80, 1996
 54. Bartels SJ, Teague GB, Drake RE, et al: Service utilization and costs associated with substance abuse among rural schizophrenic patients. *Journal of Nervous and Mental Disease* 181:227–232, 1993
 55. Dickey B, Azeni H: Persons with dual diagnosis of substance abuse and major mental illness: their excess costs of psychiatric care. *American Journal of Public Health* 86:973–977, 1996
 56. Clark RE: Family costs associated with severe mental illness and substance use. *Hospital and Community Psychiatry* 45:808–813, 1994
 57. Clark RE, Teague GB, Ricketts SK, et al: Cost-effectiveness of assertive community treatment versus standard case management for persons with co-occurring severe mental illness and substance use disorders. *Health Services Research* 33:1283–1306, 1998
 58. Jerrell JM: Cost-effective treatment for persons with dual disorders. *New Directions for Mental Health Services*, no 70:79–91, 1996
 59. O'Keefe C, Potenza DP, Mueser KT: Treatment outcomes for severely mentally ill patients on conditional discharge to community-based treatment. *Journal of Nervous and Mental Disease* 185:409–411, 1997
 60. Harris M: *Trauma Recovery and Empowerment Manual*. Edited by Anglin J. New York, Free Press, 1998
 61. Ries RK, Dyck DG: Representative payee practices of community mental health centers in Washington State. *Psychiatric Services* 48:811–814, 1997
 62. Shaner A, Roberts LJ, Eckman TA, et al: Monetary reinforcement of abstinence from cocaine among mentally ill patients with cocaine dependence. *Psychiatric Services* 48:807–810, 1997
 63. Zimmet SV, Strous RD, Burgess ES, et al: Effects of clozapine on substance use in patients with schizophrenia and schizoaffective disorders: a retrospective survey. *Journal of Clinical Psychopharmacology* 20:94–98, 2000
 64. Mueser KT, Noordsy DL, Essock S: Use of disulfiram in the treatment of patients with dual diagnosis. *American Journal of Addiction*, in press
 65. Brunette MF, Drake RE: Gender differences in patients with schizophrenia and substance abuse. *Comprehensive Psychiatry* 38:109–116, 1997
 66. Alexander MJ: Women with co-occurring addictive and mental disorders: an emerging profile of vulnerability. *American Journal of Orthopsychiatry* 66:61–70, 1996
 67. Quimby E: Homeless clients' perspectives on recovery in the Washington, DC, Dual Diagnosis Project. *Contemporary Drug Problems*, Summer 1995, pp 265–289
 68. Goodman LA, Rosenberg SD, Mueser KT, et al: Physical and sexual assault history in women with serious mental illness: prevalence, correlates, treatment, and future research directions. *Schizophrenia Bulletin* 23:685–696, 1997
 69. Mueser KT, Goodman LB, Trumbetta SL, et al: Trauma and posttraumatic stress disorder in severe mental illness. *Journal of Consulting and Clinical Psychology* 66:493–499, 1998
 70. Surgeon General's Report on Mental Health. Washington, DC, US Government Printing Office, 2000
 71. Ridgely M, Goldman H, Willenbring M: Barriers to the care of persons with dual diagnoses: organizational and financing issues. *Schizophrenia Bulletin* 16:123–132, 1990
 72. Mark T, McKusick D, King E, et al: *National Expenditures for Mental Health, Alcohol, and Other Drug Abuse Treatment*. Rockville, Md, Substance Abuse and Mental Health Services Administration, 1998
 73. Drainoni M, Bachman S: Overcoming treatment barriers to providing services for adults with dual diagnosis: three approaches. *Journal of Disability Policy Studies* 6:43–55, 1995
 74. Carey KB, Purnine DM, Maisto SM, et al: Treating substance abuse in the context of severe and persistent mental illness: clinicians' perspectives. *Journal of Substance Abuse Treatment* 19:189–198, 2000
 75. Clark RE: Family support and substance use outcomes for persons with mental illness and substance use disorders. *Schizophrenia Bulletin*, in press
 76. Mueser KT, Drake R, Wallach M: Dual diagnosis: a review of etiological theories. *Addictive Behaviors* 23:717–734, 1998
 77. *Co-occurring Psychiatric and Substance Disorders in Managed Care Systems*. Rockville, Md, Mental Health Services, 1998
 78. Hesketh B: Dilemmas in training for transfer and retention. *Applied Psychology* 46:317–386, 1997
 79. Milne D, Gorenski O, Westerman C, et al: What does it take to transfer training? *Psychiatric Rehabilitation Skills* 4:259–281, 2000
 80. Rapp CA, Poertner J: *Social Administration: A Client-Centered Approach*. White Plains, NY, Longman, 1992
 81. Mercer-McFadden C, Drake RE, Clark RE, et al: *Substance Abuse Treatment for People With Severe Mental Disorders*. Concord, NH, New Hampshire–Dartmouth Psychiatric Research Center, 1998
 82. Corrigan PW: Wanted: champions of rehabilitation for psychiatric hospitals. *American Psychologist* 40:514–521, 1995
 83. Mueser KT, Fox L: *Stagewise Family Treatment for Dual Disorders: Treatment Manual*. Concord, NH, New Hampshire–Dartmouth Psychiatric Research Center, 1998
 84. Torrey WC, Drake RE, Dixon L, et al: Implementing evidence-based practices for persons with severe mental illnesses. *Psychiatric Services* 52:45–50, 2001
 85. Shaner A, Khaka E, Roberts L, et al: Unrecognized cocaine use among schizophrenic patients. *American Journal of Psychiatry* 150:777–783, 1993
 86. Ananth J, Vandewater S, Kamal M, et al: Missed diagnosis of substance abuse in psychiatric patients. *Hospital and Community Psychiatry* 40:297–299, 1989
 87. Carey KB, Bradizza CM, Stasiewicz PR, et al: The case for enhanced addictions training in graduate programs. *Behavior Therapist* 22:27–31, 1999

The Evidence

Selected Bibliography

Literature reviews

Drake, R. E., & Brunette, M. F. (1998). Complications of severe mental illness related to alcohol and other drug use disorders. In M. Galanter (Ed.), *Recent developments in alcoholism (Vol. XIV, Consequences of alcoholism: medical, neuropsychiatric, economic, cross-cultural)*, pp. 285-299). New York: Plenum.

Summarizes research on the effects of alcohol and drug abuse on the course of severe mental illness.

Drake, R. E., Mueser, K. T., Brunette, M. F., & McHugo, G.J. (2004). A review of treatments for people with severe mental illnesses and co-occurring substance use disorders. *Psychiatric Rehabilitation Journal*, 27(4), 360-374.

- Is a comprehensive review of research on Integrated Treatment for Co-Occurring Disorders.
- Covers a wide range of research, including early demonstration programs in establishing the feasibility of Integrated Treatment in community support service settings.



Drake, R. E., Essock, S. M., Shaner, A., Carey, K. B., Minkoff, K., Kola, L., et al. (2001). Implementing dual diagnosis services for clients with severe mental illness. *Psychiatric Services*, 52(4), 469-476.

Update on research on Integrated Treatment for Co-Occurring Disorders.

Mueser, K. T., & Kavanagh, D. (2001). Treating comorbidity of alcohol problems and psychiatric disorder. In N. Heather, T. J. Peters, & T. R. Stockwell (Eds.), *International handbook of alcohol dependence and problems* (pp. 627-647). Chichester, England: Wiley.

- Provides an updated review of the epidemiology of co-occurring disorders.
- Describes integrated treatment approaches for co-occurring disorders with specific recommendations for different types of psychiatric disorders, including schizophrenia, bipolar disorder, depression, and anxiety disorders.

Mueser, K. T., Drake, R. E., & Wallach, M. A. (1998). Dual diagnosis: A review of etiological theories. *Addictive Behaviors*, 23(6), 717-734.

- Reviews the research literature on different theories accounting for the high rate of substance abuse in consumers.
- Challenges the prevailing hypothesis that high rates of substance abuse and serious mental illness could be explained by “self-medicating” distressful symptoms.
- Marshals evidence suggesting that some excess comorbidity is due to increased biological sensitivity to the effects of drugs and alcohol in consumers.

U.S. Department of Health and Human Services. (2002). *Report to Congress on the prevention and treatment of co-occurring substance abuse disorders and mental disorders*. Substance Abuse Mental Health Administration, Rockville, MD.

Includes a summary of the characteristics and needs of people with co-occurring disorders, a description of evidence-based practices and a *Five-Year Blueprint for Action*.

Historical context for Integrated Treatment for Co-Occurring Disorders

Kushner, M. G., & Mueser, K. T. (1993). Psychiatric co-morbidity with alcohol use disorders. In *Eighth special report to the U.S. Congress on alcohol and health* (NIH Pub. No. 94-3699, pp. 37-59). Rockville, MD: U.S. Department of Health and Human Services.

- Early comprehensive review of the epidemiology, correlates, and outcome of co-occurring disorders.
- Summarizes research on the negative effects of psychiatric comorbidity on the course and outcome of treatment for substance abuse.

Polcin, D. L. (1992). Issues in the treatment of dual diagnosis clients who have chronic mental illness. *Professional Psychology: Research and Practice*, 23(1), 30-37.

Describes obstacles in traditional treatment approaches to effective intervention for co-occurring disorders.

Ridgely, M. S., Goldman, H. H., & Willenbring, M. (1990). Barriers to the care of persons with dual diagnoses: Organizational and financing issues. *Schizophrenia Bulletin*, 16(1), 123-132.

Ridgely, M. S., Osher, F. C., Goldman, H. H., & Talbott, J. A. (1987). Executive summary: Chronic mentally ill young adults with substance abuse problems: A review of research, treatment, and training issues. Baltimore: University of Maryland School of Medicine, mental Health Policy Studies.

These two publications summarize problems with traditional approaches to co-occurring disorders, including administrative, clinical, and philosophical barriers to accessing intervention for both disorders.

Principles of Integrated Treatment for Co-Occurring Disorders

Minkoff, K. (1989). An integrated treatment model for dual diagnosis of psychosis and addiction. *Hospital and Community Psychiatry*, 40(10), 1031-1036.

Describes essential ingredients for an integrated service system for treating serious mental illnesses and substance use disorders.

Mueser, K. T., Drake, R. E., & Noordsy, D. L. (1998). Integrated mental health and substance abuse treatment for severe psychiatric disorders. *Practical Psychiatry and Behavioral Health*, 4, 129-139.

- Summarizes the fundamental ingredients of effective Integrated Treatment programs, including comprehensiveness, assertive outreach, assertive and protective living environment, motivation-based intervention, and long-term perspective.

- Explains the stages of treatment (engagement, persuasion, active treatment, relapse prevention), which guide practitioners in selecting interventions appropriate for consumers' level of motivation to address substance use disorders.

Ziedonis, D., & Fisher, W. (1996). Motivation-based assessment and treatment of substance abuse in patients with schizophrenia. *Directions in Psychiatry*, 16, 1-7.

Provides overview of motivation-based approach to assess and treat co-occurring disorders.

Implementation and administrative issues

Drake, R. E., Essock, S. M., Shaner, A., Carey, K. B., Minkoff, K., Kola, L., et al. (2001). Implementing dual diagnosis services for clients with severe mental illness. *Psychiatric Services*, 52(4), 469-476.

Covers issues related to implementing and disseminating Integrated Treatment for Co-Occurring Disorders.

Fox, T., Fox, L., & Drake, R. E. (1992). Developing a statewide service system for people with co-occurring severe mental illness and substance use disorders. *Innovations and Research*, 1(4), 9-13.

Describes developing integrated co-occurring disorder services in the state of New Hampshire.

Torrey, W. C., Drake, R. E., Cohen, M., Fox, L. B., Lynde, D., Gorman, P., et al. (2002). The challenge of implementing and sustaining integrated dual disorders treatment programs. *Community Mental Health Journal*, 38(6), 507-521.

State and local administrative perspectives

Fox, T. S., & Shumway, D. L. (1995). Human resource development: Training and motivating staff to work with dually diagnosed clients. In A. F. Lehman & L.B. Dixon (Eds.), *Double jeopardy: Chronic mental illness and substance use disorders* (pp. 265-276). Switzerland: Harwood Academic Publishers.

Describes how to cultivate practitioners and administrators in developing Integrated Treatment for Co-Occurring Disorders.

Mercer-McFadden, C., Drake, R. E., Clark, R. E., Verven, N., Noordsy, D. L., & Fox, T. S. (1998). *Substance abuse treatment for people with severe mental disorders: A program manager's guide*. Concord, NH: New Hampshire-Dartmouth Psychiatric Research Center.

Useful guide for program managers and anyone else with administrative responsibility for establishing and maintaining high-quality integrated programs for co-occurring disorders.

Financing and cost-effectiveness of Integrated Treatment for Co-Occurring Disorders

Clark, R. E., Teague, G. B., Ricketts, S. K., Bush, P. W., Xie, H., McGuire, T. G., et al. (1998). Cost-effectiveness of assertive community treatment versus standard case management for persons with co-occurring severe mental illness and substance use disorders. *Health Services Research*, 33(5 Pt.1), 1285-1308.

Describes cost-effectiveness analysis of study comparing Assertive Community Treatment (ACT) with standard case management for co-occurring disorders.

Clark, R. E., Ricketts, S. K., & McHugo, G. J. (1999). Legal system involvement and costs for persons in treatment for severe mental illness and substance use disorders. *Psychiatric Services*, 50, 641-647.

Addresses cost of legal system involvement in consumers with co-occurring disorders.

Dickey, B., & Azeni, H. (1996). Persons with dual diagnoses of substance abuse and major mental illness: Their excess costs of psychiatric care. *American Journal of Public Health*, 86(7), 973-977.

Documents the high cost of standard (non-integrated) treatment approaches to substance abuse in consumers.

Additional readings for program leaders and public mental health authorities

Batalden, P. B., & Stoltz, P. K. (1993). A framework for the continual improvement of health care: Building and applying professional and improvement knowledge to test changes in daily work. *The Joint Commission Journal on Quality Improvement*, 19(10), 424-447.

Gowdy, E. A., & Rapp, C. A. (1989). Managerial behavior: The common denominators of effective community based programs. *Psychosocial Rehabilitation Journal*, 13(2), 31-51.

Mercer, C. C., Mueser, K. T., & Drake, R. E. (1998) Organizational guidelines for dual disorders programs. *The Psychiatric Quarterly* 69(3), 145-168.

Nelson, E. C., Batalden, P. B., Ryer, J. C. (Eds.). (1998). *Clinical improvement action guide*. Oakbrook Terrace, Illinois: Joint Commission on Accreditation of Healthcare Organizations.

Rapp, C. A. (1993) Client-centered performance management for rehabilitation and mental health services. In R. W. Flexer & P. L. Solomon (Eds.), *Community and social support for people with severe mental disabilities*. Boston, MA: Andover. pp. 183-192.

Rapp, C. A. (1993). Client-centered performance management and the inverted hierarchy. In R. W. Flexer & P. L. Solomon (Eds.), *Community and social support for people with severe mental disabilities*. Boston, MA: Andover.

Supervisor's Tool Box. (1997). Lawrence: The University of Kansas School of Social Welfare.

Fidelity measures for Integrated Treatment for Co-Occurring Disorders

Jerrell, J. M., & Ridgely, M. S. (1999). Impact of robustness of program implementation on outcomes of clients in dual diagnosis programs. *Psychiatric Services*, 50(1), 109-112.

Documents that better substance abuse outcomes in consumers with co-occurring disorders are associated with higher program fidelity to the Integrated Treatment model.

McHugo, G. J., Drake, R. E., Teague, G. B., & Xie, H. (1999). Fidelity to assertive community treatment and client outcomes in the New Hampshire dual disorders study. *Psychiatric Services*, 50(6), 818-824.

Shows that consumers in Assertive Community Treatment (ACT) programs that implemented Integrated Treatment with high fidelity to the model had better substance abuse outcomes than those in low-fidelity programs.

Practice manuals

Center for Substance Abuse Treatment (2005). *Substance abuse treatment for persons with co-occurring disorders*. Treatment Improvement Protocol (TIP) Series 42: DHHS Publication No. (SMA) 05-3922. Rockville, MD: Substance Abuse and Mental Health Services Administration.

- Provides practical information about treating consumers with co-occurring disorders.
- Has separate chapters on treatment systems, linkages for mental health and substance abuse treatment, mood disorders, anxiety disorders, personality disorders, psychotic disorders, and pharmacology management.

Drake, R. E., Merrens, M. R., & Lynde, D. W. (2005). *Evidence-based mental health practice: A textbook*. New York: W.W. Norton.

- Introduces the concepts and approaches of evidence-based practices for treating serious mental illnesses.
- Describes the importance of research in intervention science and the evolution of evidence-based practices.
- Includes a chapter for each of five evidence-based practices and provides historical background, practice principles, and an introduction to implementation. Vignettes highlight the experiences of staff and consumers.
- Provides an excellent, readable primer for the *Evidence-Based Practices Implementation Resource Kits*.

Mueser, K. T., Noordsy, D. L., Drake, R. E., & Fox, L. (2003). *Integrated treatment for dual diagnosis: Effective intervention for severe mental illness and substance abuse*. New York: Guilford Press.

- Provides a comprehensive clinical guide for treating co-occurring disorders from which some of the material in the KIT is drawn.

- Has information on assessment, including forms and instruments.
- Describes ancillary treatment strategies, such as residential and other housing approaches, involuntary intervention, vocational rehabilitation, and psychopharmacology.
- Gives detailed guidelines and vignettes. Individual (including case management, motivational interviewing, and cognitive behavioral counseling), group (including persuasion, active treatment, social skills training, and self-help groups), and family (including individual family and multiple-family group) approaches are described.
- Includes educational handouts that cover different topics on mental illness, substance abuse, and their interactions which can be duplicated for education with consumers and family members.

Watkins, T. R., Lewellen, A., & Barrett, M. C. (2001). *Dual diagnosis: An integrated approach to treatment*. Thousand Oaks, CA: Sage Publications.

- Discusses strategies for integrating substance abuse treatment with care for mental illness.
- Has separate chapters that address different psychiatric disorders, including schizophrenia, bipolar disorder, depression, anxiety disorders, and severe personality disorders.

Substance abuse

Johns, A. (2001). Psychiatric effects of cannabis. *The British Journal of Psychiatry* 178; 116-122.

Kinney, J. (2000). *Loosening the grip: A handbook of alcohol information* (6th ed.). Boston: McGraw-Hill.

Thombs, D. L. (1999). *Introduction to addictive behaviors*. 2nd ed. The Guilford substance abuse series. New York: Guilford Press.

Weiss, R. D., & Mirin, S. M. (1987). *Cocaine*. Washington, D.C.: American Psychiatric Press.

Assessment and treatment planning

Carey, K. B., & Correia, C. J. (1998). Severe mental illness and addictions: Assessment considerations. *Addictive Behaviors*, 23(6), 735-748.

Discusses common issues practitioners face in assessing substance abuse in consumers and provides solutions to those problems.

Connors, G. J., Donovan, D. M., & DiClemente, C. C. (2001). *Substance abuse treatment and the stages of change: Selecting and planning interventions*. The Guilford substance abuse series. New York: Guilford Press.

A helpful book on treatment planning based on consumers' motivation to change their addictive behavior.

Donovan, D. M. and Marlatt, G. A (Eds.) (1988). *Assessment of addictive behaviors*. The Guilford behavioral assessment series. New York: Guilford Press.

- The introductory chapter on assessment of addictive behaviors is outstanding.
- This book also has many chapters on specific drugs and approaches that are quite good.

Drake, R. E., Rosenberg, S. D., & Mueser, K. T. (1996). Assessing substance use disorder in persons with severe mental illness. *New Directions for Mental Health Services*, 70, 3-17.

Describes many obstacles to accurate assessment of substance abuse in consumers with co-occurring disorders and strategies for overcoming these obstacles.

McHugo, G. J., Drake, R. E., Burton, H. L., & Ackerson, T. H. (1995). A scale for assessing the stage of substance abuse treatment in persons with severe mental illness. *The Journal of Nervous and Mental Disease*, 183(17), 762-767.

Contains information about assessing stages of treatment for consumers with co-occurring disorders.

Mueser, K. T., Drake, R. E., Clark, R. E., McHugo, G. J., Mercer-McFadden, C., & Ackerson, T. (1995). *Toolkit for evaluating substance abuse in persons with severe mental illness*. Cambridge, MA: Evaluation Center @ HSRI.

- Describes three practitioner-administered scales for consumers with co-occurring disorders, including the Alcohol Use Scale, the Drug Use Scale, and the Substance Abuse Treatment Scale.
- Includes software with the scales and psychometric testing for the scales.
- Has information about training practitioners on using the scales, establishing and maintaining reliability, and validity.

Noordsy, D. L., McQuade, D. V., & Mueser, K. T. (2002). Assessment considerations. In H. L. Graham, A. Copello, M. J. Birchwood, & K. T. Mueser (Eds.), *Substance misuse in psychosis: approaches to treatment and service delivery* (pp. 159-180). Chichester, England: Wiley.

- Describes principles of assessing substance abuse in consumers.
- Explains four steps of assessment: identification, classification, functional assessment and analysis, and treatment planning.
- Describes specific methods for linking assessment to treatment.

Rosenberg, S. D., Drake, R. E., Wolford, G. L., Mueser, K. T., Oxman, T. E., Vidaver, R. M., et al. (1998). The Dartmouth assessment of lifestyle instrument (DALI): A substance use disorder screen for people with severe mental illness. *The American Journal of Psychiatry*, 155(2), 232-238.

- Describes brief screening instrument (DALI) for identifying substance abuse in consumers.
- Presents data showing that DALI outperforms other screening instruments in consumers with co-occurring disorders.

Engagement

Rapp, C. A. (1998). *The strengths model: Case management with people suffering from severe and persistent mental illness*. New York: Oxford University Press.

- Describes the engagement process in consumers.
- Helpful for anyone attempting to engage consumers with co-occurring disorders in a treatment relationship.

Stages of treatment and motivational enhancement

Carey, K. B., Purnine, D. M., Maisto, S. A., Carey, M. P., & Barnes, K. L. (1999). Decisional balance regarding substance use among persons with schizophrenia. *Community Mental Health Journal*, 35(4), 289-299.

Describes using decisional balance approach to help consumers with co-occurring disorders weigh the advantages and disadvantages of continued substance use.

D’Zurilla, T. J., & Nezu, A. M. (1999). *Problem-solving therapy: A social competence approach to clinical intervention* (2nd ed.). New York: Springer Publishing Company.

Explains problem-solving therapy, which can be applied to substance abuse or mental illness problems in consumers with co-occurring disorders.

Graham, H. L., Copello, A., Birchwood, M. J., Orford, J., McGovern, D., Maslin, J., et al. (2002). Cognitive-behavioral integrated approach for psychosis and problem substance use. In H. L. Graham, A. Copello, M. J. Birchwood, & K. T. Mueser (Eds.), *Substance misuse in psychosis: approaches to treatment and service delivery* (pp. 181-206). Chichester, England: Wiley.

- Describes cognitive-behavioral approach to treating substance abuse in consumers.
- Includes numerous useful clinical examples.

Miller, W. R., & Rollnick, S. (2002). *Motivational interviewing: Preparing people for change*. (Second ed.). New York: Guilford Press.

- About stages of change and recovery from substance abuse.
- Useful for practitioners working with consumers with co-occurring disorders.

Osher, F. C., & Kofoed, L. L. (1989). Treatment of patients with psychiatric and psychoactive substance use disorders. *Hospital and Community Psychiatry*, 40, 1025-1030.

Introduces and describes the concept of stages of treatment (engagement, persuasion, active treatment, relapse prevention) that help practitioners gear treatment interventions to consumers' individual motivational states.

Rollnick, S., Mason, P., & Butler, C. (1999). *Health behavior change: A guide for practitioners*. Edinburgh: Churchill Livingstone.

- Another helpful reference on the stages of change and recovery from substance abuse.
- Describes substance abuse counseling and relapse prevention counseling.

Marlatt, G. A., & Gordon, J. R. (1985). *Relapse prevention: Maintenance strategies in the treatment of addictive behaviors*. The Guilford clinical psychology and psychotherapy series. New York: Guilford Press.

- Describes principles of substance abuse relapse prevention.
- Was written originally for work with the substance abuse population.
- Much of the book applies to consumers who have achieved sobriety and are motivated to prevent relapses of their substance abuse.

Monti, P. M., Abrams, D. B., Kadden, R. M., & Cooney, N. L. (1989). *Treating alcohol dependence: A coping skills training guide treatment manual for practitioners*. New York: Guilford Press.

- About cognitive-behavioral treatment of substance abuse.
- Provides an introduction to basic techniques that are effective for working with consumers with co-occurring disorders in the active treatment stage.

Group treatment for co-occurring disorders

Bellack, A. S., & DiClemente, C. C. (1999). Treating substance abuse among patients with schizophrenia. *Psychiatric Services*, 50(1), 75-80.

Describes social skills training approach to co-occurring disorders treatment.

Bellack, A. S., Mueser, K. T., Gingerich, S., & Agresta, J. (1997). *Social skills training for schizophrenia: A step-by-step guide*. Treatment manuals for practitioners. New York: Guilford Press.

- Addresses how to conduct social skills training groups for consumers.
- Gives specific curriculum (steps of skills) for helping consumers refuse substances and deal with substance abuse situations.

Mueser, K. T., & Noordsy, D. L. (1996). Group treatment for dually diagnosed clients. *New Directions for Mental Health Services*, 70, 33-51.

- Describes four different types of group interventions for co-occurring disorders, including educational, stage-wise (persuasion and active treatment), social skills training, and self-help groups.
- Uses brief clinical vignettes to illustrate different group treatment methods.

Noordsy, D. L., Schwab, B., Fox, L., & Drake, R. E. (1996). The role of self-help programs in the rehabilitation of persons with severe mental illness and substance use disorders. *Community Mental Health Journal*, v. 32(1), pp. 71-81.

Summarizes difficulties and solutions associated with using self-help groups, such as Alcoholics Anonymous, for consumers with co-occurring disorders.

Roberts, L. J., Shaner, A., & Eckman, T. A. (1999). *Overcoming addictions: Skills training for People with schizophrenia*. New York: W.W. Norton.

Manual for providing social skills training to consumers with co-occurring disorders.

Weiss, R. D., Greenfield, S. F., & O'Leary, G. (2002). Relapse prevention for patients with bipolar and substance use disorders. In H. L. Graham, A. Copello, M. J. Birchwood, & K. T. Mueser (Eds.), *Substance misuse in psychosis: Approaches to treatment and service delivery* (pp. 207-226). Chichester, England: Wiley.

- Describes group intervention program for bipolar disorder and substance abuse.
- Includes useful clinical examples to illustrate group treatment methods.

Self-help

Alcoholics Anonymous (2005). *The AA group: Where it all begins (rev.)*. New York: Alcoholics Anonymous.

Alcoholics Anonymous (AA) is the largest self-help organization for addiction. This book describes its history, traditions, and approach to recovery from addiction, based on the 12 Steps of AA. Available from http://www.aa.org/en_pdfs/p-16_theaagroup.pdf.

Hamilton, T., & Samples, P. (1994). *The twelve steps and dual disorders: A framework of recovery for those of us with addiction and an emotional or psychiatric illness*. Center City, MN: Hazelden.

Explains 12-Step approach to self-help substance abuse treatment in consumers.

The Dual Disorder Recovery Book: A twelve-step program for those of us with addiction and an emotional or psychiatric illness. (1993) Hazelden, Center City, Minnesota, 1993.

Discusses 12-step self-help approach to recovery for consumers with co-occurring disorders.

Trimpey, J. (1996). *Rational recovery: The new cure for substance addiction*. New York: Pocket Books.

- Rational Recovery (RR) is a self-help alternative to 12-step approaches (such as Alcoholics Anonymous).
- RR is less spirituality-oriented and more focused on helping consumers take control of their lives by accepting personal responsibility of themselves and others.

Vaillant, G. E. (1995). *The natural history of alcoholism revisited*. Cambridge, MA: Harvard University Press.

Analyzes natural pathways to recovery and explains how self-help and treatment can enhance the process.

Family treatment

Barrowclough, C. (2002). Family intervention for substance misuse in psychosis. In H. L. Graham, A. Copello, M. J. Birchwood, & K. T. Mueser (Eds.), *Substance misuse in psychosis: Approaches to treatment and service delivery* (pp. 227-243). Chichester: Wiley.

Describes family intervention approach for co-occurring disorders.

McFarlane, W. R. (2002). *Multifamily groups in the treatment of severe psychiatric disorders*. New York: Guilford Press.

Provides detailed guidance on forming and running multi-family groups for consumers and their families.

Mueser, K. T., & Fox, L. (2002). A family intervention program for dual disorders. *Community Mental Health Journal*, 38(3), 253-270.

- Describes family intervention program for co-occurring disorders that includes single-family sessions and multi-family group sessions.
- Presents pilot data from study of family program.

Mueser, K. T., & Gingerich, S. L. (in press). *Coping with schizophrenia: A guide for families* (2nd ed.). New York: Guilford Press.

- About family interventions.
- Includes a chapter on how family members can help a relative with a co-occurring disorder.

Mueser, K. T., & Glynn, S. M. (1999). *Behavioral family therapy for psychiatric disorders* (2nd ed.). Oakland, CA: New Harbinger Publications.

- Supplies a treatment manual for practitioners that describes family intervention model for mental illness, including strategies for addressing substance abuse in consumers with co-occurring disorders.
- Includes educational handouts on different psychiatric disorders, medications, and the interactions between mental illness and substance abuse.

Psychopharmacological treatment

Day, E., Georgiou, G., & Crome, I. (2002). Pharmacological management of substance misuse in psychosis. In H. L. Graham, A. Copello, M. J. Birchwood, & K. T. Mueser (Eds.), *Substance misuse in psychosis: Approaches to treatment and service delivery* (pp. 259-280). Chichester, England: Wiley.

Detailed chapter that describes pharmacologically managing substance use disorders, including stimulants, opioids, other drugs, and alcohol.

Drake, R. E., Xie, H., McHugo, G. J., & Green, A. I. (2000). The effects of clozapine on alcohol and drug use disorders among patients with schizophrenia. *Schizophrenia Bulletin*, 26(2), 441-449.

Summarizes positive effects of clozapine on alcoholism outcomes in people with schizophrenia.

Green, A. I., Zimmet, S. V., Strous, R. D., & Schildkraut, J. J. (1999). Clozapine for comorbid substance use disorder and schizophrenia: Do patients with schizophrenia have a reward-deficiency syndrome that can be ameliorated by clozapine? *Harvard Review of Psychiatry*, 6(6), 287-296.

Theoretical paper in which authors suggest that the neurobiology of schizophrenia makes consumers with this disorder more susceptible to substance abuse and more likely to benefit from clozapine.

Mueser, K. T., & Lewis, S. (2000). Treatment of substance misuse in schizophrenia. In P. Buckley & J. L. Waddington (Eds.), *Schizophrenia and mood disorders: The new drug therapies in clinical practice* (pp. 286-296). Oxford: Butterworth-Heinemann.

Brief chapter that includes recommendations for pharmacological treatment of consumers with co-occurring disorders.

Mueser, K. T., Noordsy, D. L., Fox, L., & Wolfe, R. (2003). Disulfiram treatment for alcoholism in severe mental illness. *The American Journal on Addictions*, 12(3), 242-252.

Quantitatively describes positive, long-term outcomes of 30 consumers and alcoholism treated with disulfiram (Antabuse).

Infectious diseases

Bartlett, J. G., & Finkbeiner, A. K. (1998). *The guide to living with HIV infection: Developed at the Johns Hopkins AIDS clinic*. Johns Hopkins Press Health Books. Baltimore, MD: The John Hopkins University Press.

Coping with HIV and hepatitis.

Eversen G. T., & Weinberg, H. L (1999). *Living with Hepatitis C: A Survivor's Guide*. New York: Hatherleigh Press.

Coping with HIV and hepatitis.

Razzano, L. (2002). Issues in comorbidity and HIV/AIDS. In H. L. Graham, A. Copello, M. J. Birchwood, & K. T. Mueser (Eds.), *Substance misuse in psychosis: Approaches to treatment and service delivery* (pp. 332-346). Chichester, England: Wiley.

Practical chapter on the nature of HIV/AIDS in consumers with co-occurring disorders and treatment approaches.

First-person accounts

Alverson, H., Alverson, M., & Drake, R. E. (2000). An ethnographic study of the longitudinal course of substance abuse among people with severe mental illness. *Community Mental Health Journal*, 36(6), 557-569.

Alverson, H., Alverson, M., & Drake, R. E. (2001). Social patterns of substance use among people with dual diagnoses. *Mental Health Services Research*, 3(1), 3-14.

Fox, L. (1999). Missing out on motherhood. *Psychiatric Services*, 50(2), 193-194.

Green, V. L. The resurrection and the life. *The American Journal of Orthopsychiatry*, 66(1), 12-16, 1996.

Family perspectives

Clark, R. E. (2001). Family support and substance use outcomes for persons with mental illness and substance use disorders. *Schizophrenia Bulletin*, 27(1), 93-101.

Describes family members' financial and time contributions in helping a relative with co-occurring disorders and the relationship between family assistance and improved outcomes.

Schwab, B., Clark, R. E., & Drake, R. E. (1991). An ethnographic note on clients as parents. *Psychosocial Rehabilitation Journal*, 15(2), 95-99.

Describes challenges faced by consumers with co-occurring disorders who are parents.

The Evidence

References

The following list includes the references for all citations in the KIT.

Barrowclough, C., Haddock, G., Tarrier, N., Lewis, S. W., Moring, J., O'Brien, R., et al. (2001). Randomized controlled trial of motivational interviewing, cognitive behavior therapy, and family intervention for patients with comorbid schizophrenia and substance use disorders. *The American Journal of Psychiatry*, *158*(10), 1706-1713.

Blankertz, L. E., & Cnaan, R. A. (1994). Assessing the impact of two residential programs for dually diagnosed homeless individuals. *Social Service Review*, *68*(4), 536-560.

Brunette, M., & Drake, R. E. (1998). Gender differences in homeless persons with schizophrenia and substance abuse. *Community Mental Health Journal*, *34*(6), 627-642.

Brunette, M. F., & Drake, R. E. (1997). Gender differences in patients with schizophrenia and substance abuse. *Comprehensive Psychiatry*, *38*(2), 109-116.

Brunette, M. F., Drake, R. E., Woods, M., & Hartnett, T. (2001). A comparison of long-term and short-term residential treatment programs for dual diagnosis patients. *Psychiatric Services*, *52*(4), 526-528.



- Burnam, M. A., Morton, S. C., McGlynn, E. A., Peterson, L. P., Stecher, B. M., Hayes, C., et al. (1995). An experimental evaluation of residential and nonresidential treatment for dually diagnosed homeless adults. *Journal of Addictive Diseases, 14*(4), 111-134.
- Carmichael, D. M., Tackett-Gibson, L., O'Dell, R., Menon, R., Jayasuria, B., & Jordan, J. (1998). *Texas dual diagnosis pilot project evaluation report*. Final report submitted to the Texas Commission on Alcohol and Drug Abuse and the Texas Department of Mental Health and Mental Retardation. Public Policy Research Institute, Texas A&M University.
- Drake, R. E., Essock, S. M., Shaner, A., Carey, K. B., Minkoff, K., Kola, L., et al. (2001). Implementing dual diagnosis services for clients with severe mental illness. *Psychiatric Services, 52*(4), 469-476.
- Drake, R. E., McHugo, G. J., Clark, R. E., Teague, G. B., Xie, H., Miles, K., et al. (1998a). Assertive community treatment for patients with co-occurring severe mental illness and substance use disorder: A clinical trial. *The American Journal of Orthopsychiatry, 68*(2), 201-215.
- Drake, R. E., Mercer-McFadden, C., Mueser, K. T., McHugo, G. J., & Bond, G. R. (1998b). Review of integrated mental health and substance abuse treatment for patients with dual disorders. *Schizophrenia Bulletin, 24*(4), 589-608.
- Drake, R. E., Yovetich, N. A., Bebout, R. R., Harris, M., & McHugo, G. J. (1997). Integrated treatment for dually diagnosed homeless adults. *Journal of Nervous and Mental Disease, 185*(5), 298-305.
- Franco, H., Galanter, M., Castaneda, R., & Patterson, J. (1995). Combining behavioral and self-help approaches in the inpatient management of dually diagnosed patients. *Journal of Substance Abuse Treatment, 12*(3), 227-232.
- Ganju, V. (2004). *Evidence-based practices: Responding to the challenge*. Presented at the 2004 NASMHPD Commissioner's Meeting, San Francisco, CA: June 22-24, 2004.
- Gearon, J. S., & Bellack, A. S. (1999). Women with schizophrenia and co-occurring substance use disorders: An increased risk for violent victimization and HIV. *Community Mental Health Journal, 35*(5), 401-419.
- Godley, S. H., Hoewing-Roberson, R., & Godley, M. D. (1994). *Final MISA report*. Bloomington, IL: Chestnut Health Systems/Lighthouse Institute.
- Hyde, P. S., Falls, K., Morris, J. A., & Schoenwald, S. K. (2003). *Turning knowledge into practice: A manual for behavioral health administrators and practitioners about understanding and implementing evidence-based practices*. Boston: Technical Assistance Collaborative, Inc. Available through <http://www.tacinc.org> or <http://www.acmha.org>.
- Institute of Medicine (2006). *Improving the quality of health care for mental and substance-use conditions: Quality Chasm Series*. Washington, DC: National Academy of Sciences.
- Jerrell, J. M., & Ridgely, M. S. (1995). Comparative effectiveness of three approaches to serving people with severe mental illness and substance abuse disorders. *The Journal of Nervous and Mental Disease, 183*(9), 566-576.
- Ley, A., & Jeffery, D. (2002). Cochrane review of treatment outcome studies and its implications for future developments. In H. L. Graham, A. Copello, M. J. Birchwood, & K. T. Mueser (Eds.), *Substance misuse in psychosis: Approaches to treatment and service delivery* (pp. 349-365). Chichester, England: Wiley.
- McHugo, G. J., Drake, R. E., Burton, H. L., & Ackerson, T. H. (1995). A scale for assessing the stage of substance abuse treatment in persons with severe mental illness. *The Journal of Nervous and Mental Disease, 183*(17), 762-767.

- Meisler, N., Blankertz, L., Santos, A. B., & McKay, C. (1997). Impact of assertive community treatment on homeless persons with co-occurring severe psychiatric and substance use disorders. *Community Mental Health Journal*, 33(2), 113-122.
- Mueser, K. T., Essock, S. M., Drake, R. E., Wolfe, R. S., & Frisman, L. (2001). Rural and urban differences in patients with dual diagnosis: Implications for service needs. *Schizophrenia Research*, 48(1), 93-107.
- Mueser, K. T., Yarnold, P. R., & Bellack, A. S. (1992). Diagnostic and demographic correlates of substance abuse in schizophrenia and major affective disorder. *Acta Psychiatrica Scandinavica*, 85(1), 48-55.
- Mueser, K. T., Yarnold, P. R., Rosenberg, S. D., Swett, C., Miles, K. M., & Hill, D. (2000). Substance use disorder in hospitalized severely mentally ill psychiatric patients: Prevalence, correlates, and subgroups. *Schizophrenia Bulletin*, 26(1), 179-192.
- National Advisory Mental Health Council Workgroup on Child and Adolescent Mental Health Intervention. *Development and deployment. Blueprint for change: Research on child and adolescent mental health*. Washington, DC: 2001. Available through <http://www.nimh.nih.gov>.
- New Freedom Commission on Mental Health. (2003). *Achieving the promise: Transforming mental health care in America. Final report*. DHHS Pub. No. SMA-03-3832. Rockville, MD.
- Penn, P. E., & Brooks, A. J. (1999). *Comparing substance abuse treatments for dual diagnosis: Final report*. Tucson, AZ: La Frontera Center.
- Peters, T. J., & Waterman, R. H. (1982). *In Search of excellence*. New York: Harper & Row.
- Rahav, M., Rivera, J. J., Nuttbrock, L., Ng-Mak, D., Sturz, E. L., Link, B. G., et al. (1995). Characteristics and treatment of homeless, mentally ill, chemical-abusing men. *Journal of Psychoactive Drugs*, 27(1), 93-103.
- Regier, D. A., Farmer, M. E., Rae, D. S., Locke, B. Z., Keith, S. J., Judd, L. L., et al. (1990). Comorbidity of mental disorders with alcohol and other drug abuse. Results from the Epidemiologic Catchment Area (ECA) Study. *JAMA*, 264(19), 2511-2518.
- Rosenberg, S. D., Drake, R. E., Wolford, G. L., Mueser, K. T., Oxman, T. E., Vidaver, R. M., et al. (1998). The Dartmouth assessment of lifestyle instrument (DALI): A substance use disorder screen for people with severe mental illness. *The American Journal of Psychiatry*, 155(2), 232-238.
- Rosenthal, R. N. (2002). An inpatient-based service model. In H. L. Graham, A. Copello, M. J. Birchwood, & K. T. Mueser (Eds.), *Substance misuse in psychosis: Approaches to treatment and service delivery* (pp. 136-155). Chichester, England: Wiley.
- U.S. Department of Health and Human Services. (1999). *Mental health: A report of the surgeon general*. Rockville, MD: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, and National Institutes of Health, National Institute of Mental Health.
- U.S. Department of Health and Human Services. (2001). *Mental health: Culture, race, and ethnicity. a supplement to mental health: A report of the surgeon general*. Rockville, MD: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services.



The Evidence

Acknowledgments

The materials included in the Integrated Treatment KIT were developed through the National Implementing Evidence-Based Practices Project. The Project's Coordinating Center—the New Hampshire-Dartmouth Psychiatric Research Center—in partnership with many other collaborators, including clinicians, researchers, consumers, family members, and administrators and operating under the direction of the Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, developed, evaluated, and revised these materials.

We wish to acknowledge the many people who contributed to all aspects of this project. In particular, we wish to acknowledge the contributors and consultants on the next few pages.



SAMHSA Center for Mental Health Services, Oversight Committee

Michael English

Division of Service and Systems Improvement
Rockville, Maryland

Neal B. Brown

Community Support Programs Branch
Division of Service and Systems Improvement
Rockville, Maryland

Sandra Black

Community Support Programs Branch
Division of Service and Systems Improvement
Rockville, Maryland

Crystal R. Blyler

Community Support Programs Branch
Division of Service and Systems Improvement
Rockville, Maryland

Pamela J. Fischer

Homeless Programs Branch
Division of Service and Systems Improvement
Rockville, Maryland

Sushmita Shoma Ghose

Community Support Programs Branch
Division of Service and Systems Improvement
Rockville, Maryland

Patricia Gratton

Division of Service and Systems Improvement
Rockville, Maryland

Betsy McDonel Herr

Community Support Programs Branch
Division of Service and Systems Improvement
Rockville, Maryland

Larry D. Rickards

Homeless Programs Branch
Division of Service and Systems Improvement
Rockville, Maryland

Co-Leaders

Mary Brunette

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

Robert E. Drake

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

David W. Lynde

Dartmouth Psychiatric Research Center
Concord, New Hampshire

Contributors

Stephen T. Baron

Department of Mental Health
Washington, D.C.

Gary R. Bond

Indiana University–Purdue University
Indianapolis, Indiana

John S. Caswell

West Lebanon, New Hampshire

Kevin Curdie

Concord, New Hampshire

Michael J. Cohen

National Alliance on Mental Illness (NAMI)
Concord, New Hampshire

Cathy Donahue

Calais, Vermont

Kana Enomoto

Substance Abuse and Mental Health Services
Administration
Rockville, Maryland

Carol Furlong

Harbor Homes, Inc.
Nashua, New Hampshire

Paul G. Gorman

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

Pablo R. Hernandez

Wyoming State Hospital
Evanston, Wyoming

Marta J. Hopkinson

Mental Health Services
University of Maryland–College Park
College Park, Maryland

Lenore A. Kola

Center for Evidence-Based Practice at Case
Western Reserve University
Cleveland, Ohio

Charlene LeFauve

Co-occurring and Homeless Activities Branch
Substance Abuse and Mental Health Services
Administration
Center for Substance Abuse Treatment
Rockville, Maryland

Doug Marty

The University of Kansas
Lawrence, Kansas

Gregory J. McHugo

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

Alan C. McNabb

Ascutney, Vermont

Matthew Merrens

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

Gary Morse

Community Alternatives
St. Louis, Missouri

Kim T. Mueser

Dartmouth Psychiatric Research Center
Concord, New Hampshire

Fred C. Osher

University of Maryland School of Medicine
Baltimore, Maryland

Ernest Quimby

Howard University
Washington, D.C.

Charles A. Rapp

The University of Kansas
Lawrence, Kansas

Lawrence D. Rickards

Homeless Programs Branch
Substance Abuse and Mental Health Services
Administration
Center for Mental Health Services
Rockville, Maryland



Dennis Ross

Marshfield, Vermont

Marian Scheinholtz

Community Support Programs Branch
Division of Service and Systems Improvement
Rockville, Maryland

Karin Swain

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

William Torrey

Dartmouth Medical School
Hanover, New Hampshire

Loralee West

Westbridge Community Services
Manchester, New Hampshire

Consultants to the National Implementing Evidence-Based Practices Project

Dan Adams

St. Johnsbury, Vermont

Diane C. Alden

New York State Office of Mental Health
New York, New York

Lindy Fox Amadio

Dartmouth Psychiatric Research Center
Concord, New Hampshire

Diane Asher

The University of Kansas
Lawrence, Kansas

Stephen R. Baker

University of Maryland School of Medicine
Baltimore, Maryland

Stephen T. Baron

Department of Mental Health
Washington, D.C.

Deborah R. Becker

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

Nancy L. Bolton

Cambridge, Massachusetts

Patrick E. Boyle

Case Western Reserve University
Cleveland, Ohio

Mike Brady

Adult and Child Mental Health Center
Indianapolis, Indiana

Ken Braiterman

National Alliance on Mental Illness (NAMI)
Concord, New Hampshire

Janice Braithwaite

Snow Hill, Maryland

Michael Brody

Southwest Connecticut Mental Health Center
Bridgeport, Connecticut

Mary Brunette

Dartmouth Psychiatric Research Center
Concord, New Hampshire

Sharon Bryson

Ashland, Oregon

Barbara J. Burns

Duke University School of Medicine
Durham, North Carolina

Jennifer Callaghan

The University of Kansas
School of Social Welfare
Lawrence, Kansas

Kikuko Campbell

Indiana University–Purdue University
Indianapolis, Indiana

Linda Carlson

University of Kansas
Lawrence, Kansas

Diana Chambers

Department of Health Services
Burlington, Vermont

Alice Claggett

University of Toledo College of Medicine
Toledo, Ohio

Marilyn Cloud

Department of Health and Human Services
Concord, New Hampshire

Melinda Coffman

The University of Kansas
Lawrence, Kansas

Jon Collins

Office of Mental Health and Addiction Services
Salem, Oregon

Laurie Coots

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

Judy Cox

New York State Office of Mental Health
New York, New York

Harry Cunningham

Dartmouth Psychiatric Research Center
Concord, New Hampshire

Gene Deegan

University of Kansas
Lawrence, Kansas

Natalie DeLuca

Indiana University–Purdue University
Indianapolis, Indiana

Robert E. Drake

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

Molly Finnerty

New York State Office of Mental Health
New York, New York

Laura Flint

Dartmouth Evidence-Based Practices Center
Burlington, Vermont

Vijay Ganju

National Association of State Mental Health
Program Directors Research Institute
Alexandria, Virginia

Susan Gingerich

Narberth, Pennsylvania

Phillip Glasgow

Wichita, Kansas

Howard H. Goldman

University of Maryland School of Medicine
Baltimore, Maryland

Paul G. Gorman

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

Gretchen Grappone

Concord, New Hampshire

Eileen B. Hansen

University of Maryland School of Medicine
University of Maryland, Baltimore

Kathy Hardy

Strafford, Vermont

Joyce Hedstrom

Courtland, Kansas

Lon Herman

Department of Mental Health
Columbus, Ohio

Lia Hicks

Adult and Child Mental Health Center
Indianapolis, Indiana

Debra Hrouda

Case Western Reserve University
Cleveland, Ohio



Bruce Jensen

Indiana University–Purdue University
Indianapolis, Indiana

Clark Johnson

Salem, New Hampshire

Amanda M. Jones

Indiana University–Purdue University
Indianapolis, Indiana

Joyce Jorgensen

Department of Health and Human Services
Concord, New Hampshire

Hea-Won Kim

Indiana University–Purdue University
Indianapolis, Indiana

David A. Kime

Transcendent Visions and Crazy Nation Zines
Fairless Hills, Pennsylvania

Dale Klatzker

The Providence Center
Providence, Rhode Island

Kristine Knoll

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

Bill Krenek

Department of Mental Health
Columbus, Ohio

Rick Kruszynski

Case Western Reserve University
Cleveland, Ohio

H. Stephen Leff

The Evaluation Center at the Human Services
Research Institute
Cambridge, Massachusetts

Treva E. Lichti

National Association on Mental Illness (NAMI)
Wichita, Kansas

Wilma J. Lutz

Ohio Department of Mental Health
Columbus, Ohio

Anthony D. Mancini

New York State Office of Mental Health
New York, New York

Paul Margolies

Hudson River Psychiatric Center
Poughkeepsie, New York

Tina Marshall

University of Maryland School of Medicine
Baltimore, Maryland

Ann McBride (deceased)

Oklahoma City, Oklahoma

William R. McFarlane

Maine Medical Center
Portland, Maine

Mike McKasson

Adult and Child Mental Health Center
Indianapolis, Indiana

Alan C. McNabb

Ascutney, Vermont

Meka McNeal

University of Maryland School of Medicine
Baltimore, Maryland

Ken Minkoff

ZiaLogic
Albuquerque, New Mexico

Michael W. Moore

Office of Mental Health and Addiction Services
Salem, Oregon

Roger Morin

The Center for Health Care Services
San Antonio, Texas

Lorna Moser

Indiana University–Purdue University
Indianapolis, Indiana

Kim T. Mueser

Dartmouth Psychiatric Research Center
Concord, New Hampshire

Britt J. Myrhol

New York State Office of Mental Health
New York, New York

Bill Naughton

Southeastern Mental Health Authority
Norwich, Connecticut

Nick Nichols

Department of Health
Burlington, Vermont

Bernard F. Norman

Northeast Kingdom Human Services
Newport, Vermont

Linda O'Malia

Oregon Health and Science University
Portland, Oregon

Ruth O. Ralph

University of Southern Maine
Portland, Maine

Angela L. Rollins

Indian University–Purdue University
Indianapolis, Indiana

Tony Salerno

New York State Office of Mental Health
New York, New York

Diana C. Seybolt

University of Maryland School of Medicine
Baltimore, Maryland

Patricia W. Singer

Santa Fe, New Mexico

Mary Kay Smith

University of Toledo
Toledo, Ohio

Diane Sterenbuch

Bethesda, Maryland

Bette Stewart

University of Maryland School of Medicine
Baltimore, Maryland

Steve Stone

Mental Health and Recovery Board
Ashland, Ohio

Maureen Sullivan

Department of Health and Human Services
Concord, New Hampshire

Beth Tanzman

Vermont Department of Health
Burlington, Vermont

Greg Teague

University of Southern Florida
Tampa, Florida

Boyd J. Tracy

Dartmouth Psychiatric Research Center
Lebanon, New Hampshire

Laura Van Tosh

Olympia, Washington

Joseph A. Vero

National Association on Mental Illness (NAMI)
Aurora, Ohio

Barbara L. Wieder

Case Western Reserve University
Cleveland, Ohio

Mary Woods

Westbridge Community Services
Manchester, New Hampshire



Special thanks to

The SAMHSA Center for Substance Abuse Treatment (CSAT) for partnering in reviewing the KIT and coordinating the parallel distribution of TIP 42.

The following organizations are also recognized for their generous contributions:

The Robert Wood Johnson Foundation

The John D. & Catherine T. MacArthur Foundation

West Family Foundation

Production, editorial, and graphics support

Carolyn Boccella Bagin

Center for Clear Communication, Inc.
Rockville, Maryland

Sushmita Shoma Ghose

Westat
Rockville, Maryland

Julien Hofberg

Westat
Rockville, Maryland

Glynis Jones

Westat
Rockville, Maryland

Chandria Jones

Westat
Rockville, Maryland

Tina Marshall

Gaithersburg, Maryland

Mary Anne Myers

Westat
Rockville, Maryland

Robin Ritter

Westat
Rockville, Maryland

